GRM4 Rabbit Polyclonal Antibody

CAB7284

Size:

115kDa

101kDa

Product Information

20uL, 50uL, 100uL, 200uL

83kDa/86kDa/87kDa/88kDa/



L-glutamate is the major excitatory neurotransmitter in the central nervous system and activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors, that have been divided into 3 groups on the basis of sequence homology, putative signal transduction mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5 and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3 while Group III includes GRM4, GRM6, GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic AMP cascade but differ in their agonist selectivities. Several transcript variants encoding different isoforms have been found for this gene.

Applications:

Observed MW:

Calculated MW:

WB

Reactivity:

Mouse, Rat

Antibody Information

Recommended dilutions: WB 1:500 - 1:2000

WB 1:500 - 1:2000

Source: Rabbit

Recombinant fusion protein containing a sequence corresponding to amino acids 33-290 of human GRM4 (NP_000832.1).

Isotype: IgG

Storage:

Gene ID:

2914


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Purification: Affinity purification



Uniprot Q14833 Synonyms: GRM4; GPRC1D; MGLUR4; mGlu4 Immunogen: Recombinant fusion protein containing a to amino acide 22, 200 of human GBM4

Immunogen information



Western blot analysis of extracts of various cell lines, using GRM4 antibody (CAB7284) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.