OAZ1 Rabbit Polyclonal Antibody



CAB7444

Product Information

Size:

20uL, 50uL, 100uL, 200uL

Observed MW:

27kDa

Calculated MW:

25kDa

Applications:

WB IHC

Human, Mouse, Rat

Reactivity:

Antibody Information

Recommended dilutions: WB 1:500 - 1:2000 IHC 1:50

- 1:200

Source:

Rabbit

Isotype:

IgG

Purification: Affinity purification **Protein Background**

The protein encoded by this gene belongs to the ornithine decarboxylase antizyme family, which plays a role in cell growth and proliferation by regulating intracellular polyamine levels. Expression of antizymes requires +1 ribosomal frameshifting, which is enhanced by high levels of polyamines. Antizymes in turn bind to and inhibit ornithine decarboxylase (ODC), the key enzyme in polyamine biosynthesis; thus, completing the auto-regulatory circuit. This gene encodes antizyme 1, the first member of the antizyme family, that has broad tissue distribution, and negatively regulates intracellular polyamine levels by binding to and targeting ODC for degradation, as well as inhibiting polyamine uptake. Antizyme 1 mRNA contains two potential in-frame AUGs; and studies in rat suggest that alternative use of the two translation initiation sites results in N-terminally distinct protein isoforms with different subcellular localization. Alternatively spliced transcript variants have also been noted for this gene.

Immunogen information

Gene ID:

4946

Uniprot

P54368

Synonyms:

OAZ1; AZ1; AZI; OAZ

Immunogen:

Recombinant fusion protein containing a sequence corresponding


to amino acids 1-68 of human OAZ1 (NP_004143.1).


Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.

Product Images





Western blot analysis of extracts of various cell lines, using OAZ1 antibody (CAB7444) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 60s.

Immunohistochemistry of paraffin-embedded rat kidney using OAZ1 antibody (CAB7444) at dilution of 1:100 (40x lens).