AVPR2 Rabbit Polyclonal Antibody





Product Information

Product SKU: CAB7879 **Gene ID**: 554 **Size**: 20uL, 100uL

Clone No: - Host Species: Rabbit Reactivity: Human

Additional Information

Observed MW: 46kDa **Conjugate:** Unconjugated

Calculated MW: 40kDa Isotype: IgG

Immunogen Information

Background: This gene encodes the vasopressin receptor, type 2, also known as the V2 receptor, which belongs to

the seven-transmembrane-domain G protein-coupled receptor (GPCR) superfamily, and couples to Gs thus stimulating adenylate cyclase. The subfamily that includes the V2 receptor, the V1a and V1b vasopressin receptors, the oxytocin receptor, and isotocin and mesotocin receptors in non-mammals, is well conserved, though several members signal via other G proteins. All bind similar cyclic nonapeptide hormones. The V2 receptor is expressed in the kidney tubule, predominantly in the distal convoluted tubule and collecting ducts, where its primary property is to respond to the pituitary hormone arginine vasopressin (AVP) by stimulating mechanisms that concentrate the urine and maintain water homeostasis in the organism. When the function of this gene is lost, the disease Nephrogenic Diabetes Insipidus (NDI) results. The V2 receptor is also expressed outside the kidney although its tissue localization is uncertain. When these 'extrarenal receptors' are stimulated by infusion of a V2 selective agonist (dDAVP), a variety of clotting factors are released into the bloodstream. The physiologic importance of this property is not known - its absence does not appear to be detrimental in NDI patients. The gene expression has also been described in fetal lung tissue and lung cancer associated with

alternative splicing.

Recommended Dilution: WB,1:500 - 1:2000

Synonyms: DI1; DIR; NDI; V2R; ADHR; DIR3; NDI1; AVPR2

Purifcation Method: Affinity purification

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 232-371 of human

AVPR2 (NP_000045.1).

Storage: Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.