
LPO Rabbit Polyclonal Antibody

CAB8053



| roduct Information | Protein Background |
|--------------------------|--|
| Size: | This gene encodes a member of the peroxidase family of proteins. The encoded preproproteir |
| 20uL, 50uL, 100uL, 200uL | is proteolytically processed to generate the mature enzyme. Following its secretion from salivary, mammary, and other mucosal glands, this enzyme catalyzes the generation of the |
| Observed MW: | antimicrobial substance hypothiocyanous acid. This gene is present in a gene cluster or chromosome 17. Alternative splicing results in multiple transcript variants, at least one of which |
| 70kDa | encodes a preproprotein that is proteolytically processed. |
| Calculated MW: | Immunogen information |
| 70kDa/80kDa | Gene ID: |
| Applications: | 4025 |
| WB | Uniprot |
| Reactivity: | P22079 |
| Human | Synonyms: |
| | LPO; SPO |
| Antibody Information | |
| Recommended dilutions: | Immunogen: |
| WB 1:500 - 1:2000 | Recombinant fusion protein containing a sequence corresponding to amino acids 27-280 of human LPO (NP_006142.1). |
| Source: Rabbit | |
| | Storage: |
| | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3. |
| lsotype: IgG | sourum azide, 50% giycerol, pm7.5. |

Purification: Affinity purification



Western blot analysis of extracts of various cell lines, using LPO antibody (CAB8053) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.