
ARNT2 Rabbit Polyclonal Antibody

CAB8060



Product Information Size:	Protein Background This gene encodes a member of the basic-helix-loop-helix-Per-Arnt-Sim (bHLH-PAS
5126.	superfamily of transcription factors. The encoded protein acts as a partner for several sensor
20uL, 50uL, 100uL, 200uL	proteins of the bHLH-PAS family, forming heterodimers with the sensor proteins that bind
Observed MW:	regulatory DNA sequences in genes responsive to developmental and environmental stimuli Under hypoxic conditions, the encoded protein complexes with hypoxia-inducible factor
79-85kDa	1alpha in the nucleus and this complex binds to hypoxia-responsive elements in enhancers and promoters of oxygen-responsive genes. A highly similar protein in mouse forms functiona
Calculated MW:	complexes with both aryl hydrocarbon receptors and Single-minded proteins, suggesting
77kDa/78kDa	additional roles for the encoded protein in the metabolism of xenobiotic compounds and the regulation of neurogenesis, respectively.
Applications:	Immunogen information
WB IHC	
Reactivity:	Gene ID: 9915
-	
Human, Mouse, Rat	Uniprot Q9HBZ2
Antibody Information	Synonyms:
Recommended dilutions: WB 1:500 - 1:2000 IHC 1:50 - 1:200	ARNT2; WEDAS; bHLHe1
Source:	
Rabbit	Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 440-690 of human ARNT2 (NP_055677.3).
lsotype:	
IgG	Storage:
	Storage. Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Purification: Affinity purification	



Western blot analysis of extracts of various cell lines, using ARNT2 antibody (CAB8060) at 1:1000 dilution._Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution._Lysates/proteins: 25ug per lane._Blocking buffer: 3% nonfat dry milk in TBST._Detection: ECL Enhanced Kit (CABM00021)._Exposure time: 90s.

Immunohistochemistry of paraffin-embedded human kidney using ARNT2 antibody (CAB8060) at dilution of 1:100 (40x lens).