## **UPK1B Rabbit Polyclonal Antibody**



## **CAB8157**

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

28kDa

**Calculated MW:** 

29kDa

WB

**Applications:** 

Аррисасіоны

Reactivity:

Human, Mouse

**Protein Background** 

The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded protein is found in the asymmetrical unit membrane (AUM) where it can form a complex with other transmembrane 4 superfamily proteins. It may play a role in normal bladder epithelial physiology, possibly in regulating membrane permeability of superficial umbrella cells or in stabilizing the apical membrane through AUM/cytoskeletal interactions. The use of alternate polyadenylation sites has been found for this gene.

Immunogen information

**Gene ID:** 7348

/348

Uniprot O75841

**Antibody Information** 

**Recommended dilutions:** 

WB 1:500 - 1:2000

**Synonyms:** 

UPK1B; TSPAN20; UPIB; UPK1

Source:

Immunogen:

Rabbit Recombinant fusion protein containing a sequence corresponding

to amino acids 110-230 of human UPK1B (NP $\_$ 008883.2).

Isotype:

IgG

Storage:


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.

**Purification:** 

Affinity purification

## **Product Images**



Western blot analysis of extracts of various cell lines, using UPK1B antibody (CAB8157) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.