## **CD160 Rabbit Polyclonal Antibody**



## **CAB8189**

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

20kDa

Calculated MW:

19kDa

**Applications:** 

WB

Reactivity:

Human, Mouse, Rat

**Antibody Information** 

**Recommended dilutions:** 

WB 1:500 - 1:2000

Source: Rabbit

Isotype: IgG

**Purification:** Affinity purification **Protein Background** 

CD160 is an 27 kDa glycoprotein which was initially identified with the monoclonal antibody BY55. Its expression is tightly associated with peripheral blood NK cells and CD8 T lymphocytes with cytolytic effector activity. The cDNA sequence of CD160 predicts a cysteine-rich, glycosylphosphatidylinositol-anchored protein of 181 amino acids with a single Ig-like domain weakly homologous to KIR2DL4 molecule. CD160 is expressed at the cell surface as a tightly disulfide-linked multimer. RNA blot analysis revealed CD160 mRNAs of 1.5 and 1.6 kb whose expression was highly restricted to circulating NK and T cells, spleen and small intestine. Within NK cells CD160 is expressed by CD56dimCD16+ cells whereas among circulating T cells its TCRqd expression is mainly restricted to bearing cells and TCRab+CD8brightCD95+CD56+CD28-CD27-cells. In tissues, CD160 is expressed on all intestinal intraepithelial lymphocytes. CD160 shows a broad specificity for binding to both classical and nonclassical MHC class I molecules.

Immunogen information

Gene ID:

11126

Uniprot O95971

Synonyms:

CD160; BY55; NK1; NK28

Immunogen:

Recombinant fusion protein containing a sequence corresponding


to amino acids 1-181 of human CD160 (NP\_008984.1).

Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.

## **Product Images**



Western blot analysis of extracts of various cell lines, using CD160 antibody (CAB8189) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 90s.