
Product Information

Product SKU:	CAB8189	Gene ID:	11126	Size:	20uL, 100uL
Clone No:	-	Host Species:	Rabbit	Reactivity:	Human,Mouse,Rat

Additional Information

Observed MW:	20kDa	Conjugate:	Unconjugated
Calculated MW:	20kDa	Isotype:	IgG

Immunogen Information

Background: CD160 is an 27 kDa glycoprotein which was initially identified with the monoclonal antibody BY55. Its expression is tightly associated with peripheral blood NK cells and CD8 T lymphocytes with cytolytic effector activity. The cDNA sequence of CD160 predicts a cysteine-rich, glycosylphosphatidylinositol-anchored protein of 181 amino acids with a single Ig-like domain weakly homologous to KIR2DL4 molecule. CD160 is expressed at the cell surface as a tightly disulfide-linked multimer. RNA blot analysis revealed CD160 mRNAs of 1.5 and 1.6 kb whose expression was highly restricted to circulating NK and T cells, spleen and small intestine. Within NK cells CD160 is expressed by CD56dimCD16+ cells whereas among circulating T cells its expression is mainly restricted to TCRgd bearing cells and to TCRab+CD8brightCD95+CD56+CD28-CD27-cells. In tissues, CD160 is expressed on all intestinal intraepithelial lymphocytes. CD160 shows a broad specificity for binding to both classical and nonclassical MHC class I molecules.

Recommended Dilution: WB,1:500 - 1:2000

Synonyms: NK1; BY55; NK28; CD160

Purification Method: Affinity purification

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-181 of human CD160 (NP_008984.1).

Storage: Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.