
EXOC2 Rabbit Polyclonal Antibody

CAB8235



roduct Information	Protein Background
Size:	The protein encoded by this gene is a component of the exocyst complex, a multi-protein
20uL, 50uL, 100uL, 200uL	complex essential for the polarized targeting of exocytic vesicles to specific docking sites or the plasma membrane. Though best characterized in yeast, the component proteins and the
Observed MW:	functions of the exocyst complex have been demonstrated to be highly conserved in highe eukaryotes. At least eight components of the exocyst complex, including this protein, are found
108kDa	to interact with the actin cytoskeletal remodeling and vesicle transport machinery. Thi interaction has been shown to mediate filopodia formation in fibroblasts. This protein has been
Calculated MW:	shown to interact with the Ral subfamily of GTPases and thereby mediate exocytosis by
104kDa	tethering vesicles to the plasma membrane. Alternative splicing results in multiple transcrip variants.
Applications:	Immunogen information
WB IF	-
Reactivity:	Gene ID: 55770
Human, Mouse, Rat	Uniprot
	Q96KP1
Antibody Information	<u>Companyana</u>
Recommended dilutions: WB 1:500 - 1:2000 IF 1:50 - 1:200	Synonyms: EXOC2; SEC5; SEC5L1; Sec5p
Source:	
Rabbit	Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-220 of human EXOC2 (NP_060773.3).
lsotype:	
lgG	Storage:
	Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Affinity purification



Western blot analysis of extracts of various cell lines, using EXOC2 antibody (CAB8235) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 30s.

Immunofluorescence analysis of U2OS cells using EXOC2 antibody (CAB8235) at dilution of 1:100. Blue: DAPI for nuclear staining.