## **PTPN18 Rabbit Polyclonal Antibody**



## **CAB8487**

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

60kDa

Calculated MW:

38kDa/50kDa

**Applications:** 

WB

Reactivity:

Human

**Protein Background** 

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, the mitotic cycle, and oncogenic transformation. This PTP contains a PEST motif, which often serves as a protein-protein interaction domain, and may be related to protein intracellular half-live. This protein can differentially dephosphorylate autophosphorylated tyrosine kinases that are overexpressed in tumor tissues, and it appears to regulate HER2, a member of the epidermal growth factor receptor family of receptor tyrosine kinases. Two transcript variants encoding different isoforms have been found for this gene.

Immunogen information

Gene ID: 26469

Uniprot Q99952

Synonyms: PTPN18; BDP1; PTP-HSCF

**Antibody Information** 

**Recommended dilutions:** 

WB 1:500 - 1:2000

Immunogen:

Recombinant fusion protein containing a sequence corresponding Source:

to amino acids 1-210 of human PTPN18 (NP\_001135842.1). Rabbit

Storage: Isotype:


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% IgG

sodium azide, 50% glycerol, pH7.3.

**Purification:** 

Affinity purification

## **Product Images**



Western blot analysis of extracts of various cell lines, using PTPN18 antibody (CAB8487) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 1s.