## **DUOX1 Rabbit Polyclonal Antibody**



## **CAB8583**

**Product Information** 

Size:

20uL, 50uL, 100uL, 200uL

**Observed MW:** 

180kDa

Calculated MW:

137kDa/177kDa

**Applications:** 

Reactivity:

WB

Mouse, Rat

**Protein Background** 

The protein encoded by this gene is a glycoprotein and a member of the NADPH oxidase family. The synthesis of thyroid hormone is catalyzed by a protein complex located at the apical membrane of thyroid follicular cells. This complex contains an iodide transporter, thyroperoxidase, and a peroxide generating system that includes proteins encoded by this gene and the similar DUOX2 gene. This protein is known as dual oxidase because it has both a peroxidase homology domain and a gp91phox domain. This protein generates hydrogen peroxide and thereby plays a role in the activity of thyroid peroxidase, lactoperoxidase, and in lactoperoxidase-mediated antimicrobial defense at mucosal surfaces. Two alternatively spliced transcript variants encoding the same protein have been described for this gene.

Immunogen information

Gene ID: 53905

Uniprot Q9NRD9

**Antibody Information** 

**Recommended dilutions:** 

WB 1:500 - 1:2000

Synonyms:

DUOX1; LNOX1; NOXEF1; THOX1

Immunogen: Source:

Recombinant fusion protein containing a sequence corresponding

to amino acids 1312-1551 of human DUOX1 (NP\_059130.2).

Isotype:

Rabbit

Storage: IgG


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.

**Purification:** 

Affinity purification

## **Product Images**



Western blot analysis of extracts of various cell lines, using DUOX1 antibody (CAB8583) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (CABM00021). Exposure time: 60s.