
## Phospho-mTOR-S2448 Rabbit Polyclonal Antibody

CABP0094



Product Information Size:	<b>Protein Background</b> The protein encoded by this gene belongs to a family of phosphatidylinositol kinase-related
20uL, 50uL, 100uL, 200uL	kinases. These kinases mediate cellular responses to stresses such as DNA damage and nutrient deprivation. This protein acts as the target for the cell-cycle arrest and immunosuppressive
Observed MW:	effects of the FKBP12-rapamycin complex. The ANGPTL7 gene is located in an intron of th gene.
289kDa	
Calculated MW:	Immunogen information
288kDa	Gene ID: 2475
Applications:	
WB IF	Uniprot P42345
Reactivity:	
Human, Mouse, Rat	<b>Synonyms:</b> FRAP; FRAP1; FRAP2; RAFT1; RAPT1; SKS; mTOR; MTOR
Antibody Information	
<b>Recommended dilutions:</b> WB 1:500 - 1:2000 IF 1:50 - 1:200	<b>Immunogen:</b> A synthetic phosphorylated peptide around S2448 of human mTOR (NP_004949.1).
Source:	
Rabbit	<b>Storage:</b> Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
lsotype:	
lgG	

**Purification:** Affinity purification



Western blot analysis of extracts of NIH/3T3 and C6 cells, using Phospho-MTOR-S2448 antibody (CABP0094) at 1:2000 dilution. NIH/3T3 cells were treated by Insulin (100nM) for 10 minutes after serum-starvation overnight. C6 cells were treated by IGF-1 (50 ng/mL) for 30 minutes after serum-starvation overnight. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% BSA. Detection: ECL Basic Kit (CABM00020). Exposure time: 30S.

Immunofluorescence analysis of HeLa cells using Phospho-MTOR-S2448 antibody (CABP0094) at dilution of 1:100. Blue: DAPI for nuclear staining.