Phospho-PRKCQ-S676 Rabbit Polyclonal **Antibody**





Product Information

Size:

50uL, 100uL, 200uL

Observed MW:

82kDa

Calculated MW:

67kDa/74kDa/81kDa

Applications:

WB IHC IF

Reactivity:

Human, Mouse, Rat

Immunogen information

the transcription factors.

Protein Background

Gene ID: 5588

Uniprot Q04759

Antibody Information

Recommended dilutions:

WB 1:500 - 1:2000 IHC 1:50 - 1:100 IF 1:100 - 1:200

Source:

Rabbit

Isotype:

IgG

Synonyms:

PRKCQ; PRKCT; nPKC-theta

Immunogen:

A phospho specific peptide corresponding to residues surrounding

Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be

activated by calcium and the second messenger diacylglycerol. PKC family members

phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a

class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role. The protein encoded by this gene is one of the PKC family

members. It is a calcium-independent and phospholipid-dependent protein kinase. This kinase

is important for T-cell activation. It is required for the activation of the transcription factors NFkappaB and AP-1, and may link the T cell receptor (TCR) signaling complex to the activation of

S676 of human PRKCQ

Storage:


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.


Purification:

Affinity purification


Product Images



Western blot analysis of extracts from PC12 cells using Phospho-PRKCQ-S676 antibody (CABP0260). Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% BSA.



Immunohistochemistry of paraffin-embedded human lung carcinoma using Phospho-PRKCQ-S676 antibody (CABP0260).



Immunofluorescence analysis of methanol-fixed HeLa cells using Phospho-PRKCQ-S676 antibody (CABP0260).