# Phospho-ERBB3-Y1328 Rabbit Polyclonal Antibody





#### **Product Information**

Size:

50uL, 100uL, 200uL

**Observed MW:** 

Refer to figures

Calculated MW:

20kDa/36kDa/77kDa/141kDa /148kDa

**Applications:** 

WB IHC IF

Reactivity:

Human, Mouse, Rat

# **Protein Background**

This gene encodes a member of the epidermal growth factor receptor (EGFR) family of receptor tyrosine kinases. This membrane-bound protein has a neuregulin binding domain but not an active kinase domain. It therefore can bind this ligand but not convey the signal into the cell through protein phosphorylation. However, it does form heterodimers with other EGF receptor family members which do have kinase activity. Heterodimerization leads to the activation of pathways which lead to cell proliferation or differentiation. Amplification of this gene and/or overexpression of its protein have been reported in numerous cancers, including prostate, bladder, and breast tumors. Alternate transcriptional splice variants encoding different isoforms have been characterized. One isoform lacks the intermembrane region and is secreted outside the cell. This form acts to modulate the activity of the membrane-bound form. Additional splice variants have also been reported, but they have not been thoroughly characterized.

# Immunogen information

Gene ID:

2065

Uniprot P21860

## **Antibody Information**

## **Recommended dilutions:**

WB 1:500 - 1:1000 IHC 1:50 - 1:100 IF 1:100 - 1:200

Source:

Rabbit

Isotype:

IgG

#### Synonyms:

ERBB3; ErbB-3; HER3; LCCS2; MDA-BF-1; c-erbB-3; c-erbB3; erbB3-S; p180-ErbB3; p45-sErbB3; p85-sErbB3

#### Immunogen:

A phospho specific peptide corresponding to residues surrounding Y1328 of human ERBB3

### Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

# **Purification:**

Affinity purification