## Phospho-NFKB1-S907 Rabbit Polyclonal Antibody





**Product Information** 

Size:

50uL, 100uL, 200uL

**Observed MW:** 

118kDa

Calculated MW:

85kDa/105kDa

**Applications:** 

WB

Reactivity:

Human

Immunogen information

is proteolytically processed.

**Protein Background** 

Gene ID:

4790

Uniprot P19838

**Antibody Information** 

**Recommended dilutions:** 

WB 1:500 - 1:2000

Source:

Rabbit

Isotype:

**Purification:** Affinity purification

IgG

Synonyms:

NFKB1; CVID12; EBP-1; KBF1; NF-kB1; NF-kappa-B; NF-kappaB;

This gene encodes a 105 kD protein which can undergo cotranslational processing by the 26S

proteasome to produce a 50 kD protein. The 105 kD protein is a Rel protein-specific

transcription inhibitor and the 50 kD protein is a DNA binding subunit of the NF-kappa-B (NFKB) protein complex. NFKB is a transcription regulator that is activated by various intra- and

extra-cellular stimuli such as cytokines, oxidant-free radicals, ultraviolet irradiation, and bacterial or viral products. Activated NFKB translocates into the nucleus and stimulates the

expression of genes involved in a wide variety of biological functions. Inappropriate activation

of NFKB has been associated with a number of inflammatory diseases while persistent inhibition of NFKB leads to inappropriate immune cell development or delayed cell growth. Alternative

splicing results in multiple transcript variants encoding different isoforms, at least one of which

NFKB-p105; NFKB-p50; NFkappaB; p105; p50

Immunogen:

A phospho specific peptide corresponding to residues surrounding


S907 of human NFKB1

Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.

## **Product Images**



Western blot analysis of extract from HeLa cells, using Phospho-NFKB1-S907 antibody (CABP0416). Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% BSA.