

Phospho-NF- κ B2-S870 Rabbit Polyclonal Antibody

CABP0419



Product Information

Size:

50uL, 100uL, 200uL

Observed MW:

120kDa

Calculated MW:

47kDa/96kDa

Applications:

WB

Reactivity:

Human

Protein Background

This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NF κ B). The NF κ B complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protein encoded by this gene can function as both a transcriptional activator or repressor depending on its dimerization partner. The p100 full-length protein is co-translationally processed into a p52 active form. Chromosomal rearrangements and translocations of this locus have been observed in B cell lymphomas, some of which may result in the formation of fusion proteins. There is a pseudogene for this gene on chromosome 18. Alternative splicing results in multiple transcript variants.

Immunogen information

Gene ID:

4791

Uniprot

Q00653

Antibody Information

Recommended dilutions:

WB 1:500 - 1:2000

Source:

Rabbit

Isotype:

IgG

Purification:

Affinity purification

Synonyms:

CVID10; H2TF1; LYT-10; LYT10; NF- κ B2; p100; p49/p100; p52


Immunogen:

A phospho specific peptide corresponding to residues surrounding S870 of human NF- κ B2

Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Product Images



Western blot analysis of extract from MDA-MB-435 cells, using Phospho-NF-κB2-S870 antibody (CABP0419). Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% BSA.