Phospho-MAP2K4-S257/T261 Rabbit Polyclonal Antibody



CABP0541

Product Information

Size:

20uL, 50uL, 100uL, 200uL

Observed MW:

40kDa

Calculated MW:

44kDa/45kDa

Applications:

лррпси попъ

Reactivity:

WB

Human, Mouse, Rat

Protein Background

This gene encodes a member of the mitogen-activated protein kinase (MAPK) family. Members of this family act as an integration point for multiple biochemical signals and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation, and development. They form a three-tiered signaling module composed of MAPKKKs, MAPKKs, and MAPKs. This protein is phosphorylated at serine and threonine residues by MAPKKKs and subsequently phosphorylates downstream MAPK targets at threonine and tyrosine residues. A similar protein in mouse has been reported to play a role in liver organogenesis. A pseudogene of this gene is located on the long arm of chromosome X. Alternative splicing results in multiple transcript variants.

Immunogen information

Gene ID: 6416

Uniprot P45985

Antibody Information

Recommended dilutions:

WB 1:500 - 1:2000

Synonyms:

MAP2K4; JNKK; JNKK1; MAPKK4; MEK4; MKK4; PRKMK4; SAPKK-1;

SAPKK1; SEK1; SERK1; SKK1

Source:

Rabbit

Immunogen: A synthetic phosphorylated peptide around S257 & T261 of

human MAP2K4 (NP_003001.1).

Isotype:

lgG Storage:


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02%

sodium azide, 50% glycerol, pH7.3.

Purification:

Affinity purification

Product Images



Western blot analysis of extracts of NIH/3T3 cells, using Phospho-MAP2K4-S257/T261 pAb (CABP0541) at 1:1000 dilution or MAP2K4 antibody (CAB14781).NIH/3T3 cells were treated by UV at room temperature for 15-30 minutes. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% BSA. Detection: ECL Basic Kit (CABM00020). Exposure time: 1s.