## Phospho-CSF1R-Y699 Rabbit Polyclonal Antibody

## CABP0607



Product Information Size:	<b>Protein Background</b> The protein encoded by this gene is the receptor for colony stimulating factor 1, a cytokine
20uL, 50uL, 100uL, 200uL	which controls the production, differentiation, and function of macrophages. This receptor mediates most if not all of the biological effects of this cytokine. Ligand binding activates the
Observed MW:	receptor kinase through a process of oligomerization and transphosphorylation. The encoded protein is a tyrosine kinase transmembrane receptor and member of the CSF1/PDGF receptor
Refer to figures	family of tyrosine-protein kinases. Mutations in this gene have been associated with a predisposition to myeloid malignancy. The first intron of this gene contains a transcriptionally
Calculated MW:	inactive ribosomal protein L7 processed pseudogene oriented in the opposite direction.
33kDa/107kDa	Alternative splicing results in multiple transcript variants. Expression of a splice variant from an LTR promoter has been found in Hodgkin lymphoma (HL), HL cell lines and anaplastic large cell
Applications:	lymphoma.
WB	Immunogen information
Reactivity:	Gene ID:
Human	1436
	Uniprot P07333
Antibody Information	
<b>Recommended dilutions:</b> WB 1:500 - 1:2000	<b>Synonyms:</b> CSF1R; C-FMS; CD115; CSF-1R; CSFR; FIM2; FMS; HDLS; M-CSF-R; BANDDOS
<b>Source:</b> Rabbit	
<b>Isotype:</b> IgG	<b>Immunogen:</b> A synthetic phosphorylated peptide around Y699 of human CSF1R (NP_005202.2).
<b>Purification:</b> Affinity purification	<b>Storage:</b> Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.