

# Phospho-VIM-S39 Rabbit Polyclonal Antibody



CABP0806

## Product Information

### Size:

20uL, 50uL, 100uL, 200uL

### Observed MW:

57KDa

### Calculated MW:

53kDa

### Applications:

WB IHC

### Reactivity:

Human, Mouse, Rat

## Antibody Information

### Recommended dilutions:

WB 1:500 - 1:2000 IHC 1:50  
- 1:100

### Source:

Rabbit

### Isotype:

IgG

### Purification:

Affinity purification

## Protein Background

This gene encodes a member of the intermediate filament family. Intermediate filaments, along with microtubules and actin microfilaments, make up the cytoskeleton. The protein encoded by this gene is responsible for maintaining cell shape, integrity of the cytoplasm, and stabilizing cytoskeletal interactions. It is also involved in the immune response, and controls the transport of low-density lipoprotein (LDL)-derived cholesterol from a lysosome to the site of esterification. It functions as an organizer of a number of critical proteins involved in attachment, migration, and cell signaling. Mutations in this gene causes a dominant, pulverulent cataract.

## Immunogen information

### Gene ID:

7431

### Uniprot

P08670

### Synonyms:

CTRCT30; HEL113; Vimentin; VIM; vimentin




### Immunogen:

A synthetic phosphorylated peptide around S39 of human VIM (NP\_003371.2).

### Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

## Product Images



Western blot analysis of extracts of NIH/3T3 cells, using Phospho-VIM-S39 antibody (CABP0806) at 1:1000 dilution. NIH/3T3 cells were treated by Paclitaxel (100 nM/ml) at 37°C for 20 hours. NIH/3T3 cells were treated by Hydroxyurea (4 mM) at 37°C for 20 hours. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00071). Exposure time: 1s.

Western blot analysis of extracts of HeLa cells, using Phospho-VIM-S39 antibody (CABP0806) at 1:1000 dilution. HeLa cells were treated by Paclitaxel (100 nM/ml) at 37°C for 20 hours. HeLa cells were treated by Hydroxyurea (4 mM) at 37°C for 20 hours. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (CABM00020). Exposure time: 180s.

Immunohistochemistry of paraffin-embedded human appendix using Phospho-VIM-S39 antibody (CABP0806) at dilution of 1:100 (40x lens).