

Phospho-c-Myc-T58/S62 Rabbit Monoclonal Antibody

CABP0988



Product Information

Size:

20uL, 50uL, 100uL

Observed MW:

51kDa

Calculated MW:

57kDa

Applications:

WB

Reactivity:

Human, Rat

Protein Background

The protein encoded by this gene is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma. There is evidence to show that alternative translation initiations from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site result in the production of two isoforms with distinct N-termini. The synthesis of non-AUG initiated protein is suppressed in Burkitt's lymphomas, suggesting its importance in the normal function of this gene. [provided by RefSeq, Jul 2008]

Immunogen information

Gene ID:

4609

Uniprot

P01106

Antibody Information

Recommended dilutions:

WB 1:500 - 1:2000

Source:

Rabbit

Isotype:

IgG

Purification:

Affinity purification

Synonyms:

MRTL; MYCC; bHLHe39; c-Myc


Immunogen:

A phospho specific peptide corresponding to residues surrounding T58/S62 of human c-Myc

Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Product Images



Western blot - Phospho-c-Myc-T58/S62 Rabbit mAb (CABP0988)