

Phospho-RPA2-S4/S8 Rabbit Polyclonal Antibody



CABP1102

Product Information

Size:

20uL, 50uL, 100uL, 200uL

Observed MW:

Refer to figures

Calculated MW:

29kDa/30kDa/38kDa

Applications:

WB

Reactivity:

Human, Mouse, Rat

Protein Background

As part of the heterotrimeric replication protein A complex (RPA/RP-A), binds and stabilizes single-stranded DNA intermediates, that form during DNA replication or upon DNA stress. It prevents their reannealing and in parallel, recruits and activates different proteins and complexes involved in DNA metabolism. Thereby, it plays an essential role both in DNA replication and the cellular response to DNA damage. In the cellular response to DNA damage, the RPA complex controls DNA repair and DNA damage checkpoint activation. Through recruitment of ATRIP activates the ATR kinase a master regulator of the DNA damage response. It is required for the recruitment of the DNA double-strand break repair factors RAD51 and RAD52 to chromatin in response to DNA damage. Also recruits to sites of DNA damage proteins like XPA and XPG that are involved in nucleotide excision repair and is required for this mechanism of DNA repair. Plays also a role in base excision repair (BER) probably through interaction with UNG. Also recruits SMARCAL1/HARP, which is involved in replication fork restart, to sites of DNA damage. May also play a role in telomere maintenance.

Immunogen information

Gene ID:

6118

Antibody Information

Recommended dilutions:

WB 1:500 - 1:2000

Source:

Rabbit

Uniprot

P15927

Synonyms:

REPA2; RP-A p32; RP-A p34; RPA32; RPA2; RP-Ap32; RP-Ap34

Isotype:

IgG

Immunogen:

A phospho specific peptide corresponding to residues surrounding S4/S8 of human RPA2.

Purification:

Affinity purification

Storage:

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Product Images
