

Anti-Mouse CD3ε [145-2C11] In Vivo Antibody - Ultra Low Endotoxin

Product Information

| Product SKU: IVMB0059 | |
|--|----|
| Size: 1mg, 5mg, 25mg, 50mg, 100mg | |
| Concentration: ≥ 5.0 mg/ml | Α. |
| Isotype: Armenian Hamster IgG | |
| Host: Armenian Hamster Clone: 145-2C11 | |
| Category: Monoclonal Antibody | 3 |
| Reactivity: Mouse | |
| Synonyms: | |

Specificity:

CD3, T3

Clone 145-2C11 recognizes an epitope on mouse CD3ε.

Formulation:

This monoclonal antibody is aseptically packaged and formulated in 0.01 M phosphate buffered saline (150 mM NaCl) PBS pH 7.2 - 7.4 with no carrier protein, potassium, calcium or preservatives added.



Endotoxin Level:

≤ 0.5 EU/mg as determined by the LAL method

Purity:

≥98% monomer by analytical SEC >95% by SDS Page

Immunogen:

H-2Kb-specific mouse cytotoxic T lymphocyte clone BM10-37

Storage and Stability:

Functional grade preclinical antibodies may be stored sterile as received at 2-8°C for up to one month. For longer term storage, aseptically aliquot in working volumes without diluting and store at -80°C. Avoid Repeated Freeze Thaw Cycles.

Product Preparation:

Functional grade preclinical antibodies are manufactured in an animal free facility using only In vitro protein free cell culture techniques and are purified by a multi-step process including the use of protein A or G to assure extremely low levels of endotoxins, leachable protein A or aggregates.

Applications

Act, B, CODEX®, CyTOF®, Depletion, FA, FC, ICC, IF, IHC FF, WB

Applications & Recommended Usage:

FC The suggested concentration for this 145-2C11 antibody for staining cells in flow cytometry is \leq 1.0 µg per 106 cells in a volume of 100 µl. Titration of the reagent is recommended for optimal performance for each application.