

Anti-Human CD11c In Vivo Antibody - Low Endotoxin

Product Information

Product SKU:

IVMB0109

Size:

1mg, 5mg, 25mg, 50mg, 100mg

Concentration:

≥ 5.0 mg/ml

Isotype:

Mouse IgG1 κ

Host:

Mouse

Clone:

3.9

Category:

Monoclonal Antibody

Reactivity:

Human

Synonyms:

Integrin αX subunit, CR4, p150, ITGAX

Specificity:

Clone 3.9 recognizes the α-chain (CD11c) of the CD11c/CD18 complex. It is specific for the I domain of CD11c. Clone 3.9 binds the activated form of CD11c and partially blocks the binding of CD11c with ICAM-4.

Formulation:



This monoclonal antibody is aseptically packaged and formulated in 0.01 M phosphate buffered saline (150 mM NaCl) PBS pH 7.2 - 7.4 with no carrier protein, potassium, calcium or preservatives added.

Endotoxin Level:

< 1.0 EU/mg as determined by the LAL method

Purity:

≥95% monomer by analytical SEC · >95% by SDS Page

Immunogen:

Rheumatoid synovial fluid cells and fibronectin purified human monocytes

Storage and Stability:

Functional grade preclinical antibodies may be stored sterile as received at 2-8°C for up to one month. For longer term storage, aseptically aliquot in working volumes without diluting and store at -80°C. Avoid Repeated Freeze Thaw Cycles.

Product Preparation:

Functional grade preclinical antibodies are manufactured in an animal free facility using only In vitro protein free cell culture techniques and are purified by a multi-step process including the use of protein A or G to assure extremely low levels of endotoxins, leachable protein A or aggregates.

Applications

B, CyTOF®, FA, FC, WB

Applications & Recommended Usage:

FC The suggested concentration for this 3.9 antibody for staining cells in flow cytometry is ≤ 2.0 µg per 10⁶ cells in a volume of 100 µl or 100µl of whole blood. Titration of the reagent is recommended for optimal performance for each application.>