

# Anti-Human TCR Vα24-Jα18 (iNKT cell) In Vivo Antibody - Low Endotoxin

# **Product Information**

Product SKU: IVMB0136

Size: 1mg, 5mg, 25mg, 50mg, 100mg

**Concentration:** ≥ 5.0 mg/ml

**Isotype:** Mouse IgG1 κ

Host: Mouse

Clone: 6B11

Category: Monoclonal Antibody

**Reactivity:** Human

**Synonyms:** TCR Vα24-Jα18, TCR Vα24-JαQ, invariant NKT cell, iNKT

# Specificity:

Clone 6B11 recognizes the invariant CDR3 region of the human V $\alpha$ 24-J $\alpha$ 18 TCR.

# Formulation:

This monoclonal antibody is aseptically packaged and formulated in 0.01 M phosphate buffered saline (150 mM NaCl) PBS pH 7.2 - 7.4 with no carrier protein, potassium, calcium or preservatives added.

Copyright © 2023 Assay Genie info@assaygenie.com www.assaygenie.com



#### Endotoxin Level:

< 1.0 EU/mg as determined by the LAL method

#### Purity:

≥95% monomer by analytical SEC · >95% by SDS Page

#### Immunogen:

#### Storage and Stability:

Functional grade preclinical antibodies may be stored sterile as received at 2-8°C for up to one month. For longer term storage, aseptically aliquot in working volumes without diluting and store at -80°C. Avoid Repeated Freeze Thaw Cycles.

#### **Product Preparation:**

Functional grade preclinical antibodies are manufactured in an animal free facility using only In vitro protein free cell culture techniques and are purified by a multi-step process including the use of protein A or G to assure extremely low levels of endotoxins, leachable protein A or aggregates.

#### Applications

CyTOF®, FC

# Applications & Recommended Usage:

FC The suggested concentration for this TCR V $\alpha$ 24-J $\alpha$ 18 antibody for staining cells in flow cytometry is  $\leq 1.0 \ \mu$ g per 106 cells in a volume of 100  $\mu$ l or 100 $\mu$ l of whole blood. Titration of the reagent is recommended for optimal performance for each application.