## **MAP2K6 Antibody**

# AssayGenie 🗳

#### **PACO14679**

Reactivity:

Rabbit

#### **Product Information**

Size: Protein Background:

50ul This gene encodes a member of the dual specificity protein kinase family, which functions as a mitogen-activated protein (MAP) kinase kinase. MAP kinases, also known

as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This protein phosphorylates and activates p38 MAP kinase in

Human, Mouse biochemical signals. This protein phosphorylates and activates p38 MAP kinase in response to inflammatory cytokines or environmental stress. As an essential component of p38 MAP kinase mediated signal transduction pathway this gape is involved in many

of p38 MAP kinase mediated signal transduction pathway, this gene is involved in many

cellular processes such as stress induced cell cycle arrest, transcription activation and

apoptosis.

Isotype: Gene ID:

IgG MAP2K6

Applications: Uniprot

ELISA, IHC P52564

Recommended dilutions: Synonyms:

ELISA:1:2000-1:5000, IHC:1:25-1:100 mitogen-activated protein kinase kinase 6


Immunogen:


Fusion protein of human MAP2K6.

Storage:

-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

### **Product Images**





The image on the left is immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using PACO14679(MAP2K6 Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification: x—200).

The image on the left is immunohistochemistry of paraffin-embedded Human lung cancer tissue using PACO14679(MAP2K6 Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification: x—200).