

Product Information

Size:

50ul

Reactivity:

Human, Mouse

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, WB, IHC

Recommended dilutions:

ELISA:1:1000-1:2000, WB:1:200-1:1000,
IHC:1:15-1:50

Protein Background:

The OCT1 transcription factor was among the first identified members of the POU transcription factor family. Members of this family contain the POU domain, a 160-amino acid, region necessary for DNA binding to the octameric sequence ATGCAAAT.

Gene ID:

POU2F1

Uniprot

P14859

Synonyms:

POU class 2 homeobox 1


Immunogen:

Fusion protein of human POU2F1.


Storage:

-20°C; C, pH7.4 PBS, 0.05% NaN₃, 40% Glycerol

Product Images



Gel: 8%SDS-PAGE, Lysate: 40 μ g, Lane: 231 cells, Primary antibody: PACO14798(POU2F1 Antibody) at dilution 1/400, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 50 seconds.



The image on the left is immunohistochemistry of paraffin-embedded Human lung cancer tissue using PACO14798(POU2F1 Antibody) at dilution 1/25, on the right is treated with fusion protein. (Original magnification: x—200).