
F3 Antibody

PACO15089



Product Information	
Size:	Protein Background:
50ul	This gene encodes coagulation factor III which is a cell surface glycoprotein. This factor
Reactivity:	enables cells to initiate the blood coagulation cascades, and it functions as the high- affinity receptor for the coagulation factor VII. The resulting complex provides a
Human	catalytic event that is responsible for initiation of the coagulation protease cascades by specific limited proteolysis. Unlike the other cofactors of these protease cascades, which
Source:	circulate as nonfunctional precursors, this factor is a potent initiator that is fully
Rabbit	functional when expressed on cell surfaces. There are 3 distinct domains of this factor: extracellular, transmembrane, and cytoplasmic. This protein is the only one in the
lsotype:	coagulation pathway for which a congenital deficiency has not been described. Alternate splicing results in multiple transcript variants.
lgG	Gene ID:
Applications:	F3
ELISA, IHC	Uniprot
Recommended dilutions:	P13726
ELISA:1:2000-1:5000, IHC:1:25-1:100	Synonyms:
	Coagulation factor III (thromboplastin, tissue factor)
	Immunogen:
	Fusion protein of human F3.
	Storage:
	-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol



The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using PACO15089(F3 Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification: x—200).

The image on the left is immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using PACO15089(F3 Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification: x—200).