ATP2C1 Antibody



PACO15839

Reactivity:

Human, Mouse, Rat

Product Information

Size: Protein Background:

50ul The protein encoded by this gene belongs to the family of P-type cation transport

ATPases. This magnesium-dependent enzyme catalyzes the hydrolysis of ATP coupled with the transport of calcium ions. Defects in this gene cause Hailey-Hailey disease, an

autosomal dominant disorder. Alternatively spliced transcript variants encoding

different isoforms have been identified.

Source: Gene ID:

Rabbit ATP2C1

Isotype: Uniprot

IgG P98194

Applications: Synonyms:

ELISA, IHC

ATPase, Ca++ transporting, type 2C, member 1


Recommended dilutions: Immunogen:


ELISA:1:1000-1:5000, IHC:1:50-1:200 Fusion protein of human ATP2C1.

Storage:

-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product Images





The image on the left is immunohistochemistry of paraffin-embedded Human colon cancer tissue using PACO15839(ATP2C1 Antibody) at dilution 1/60, on the right is treated with fusion protein. (Original magnification: x—200).

The image on the left is immunohistochemistry of paraffin-embedded Human gastic cancer tissue using PACO15839(ATP2C1 Antibody) at dilution 1/60, on the right is treated with fusion protein. (Original magnification: x—200).