

HSD17B6 Antibody



PACO16495

Product Information

Size:

50ul

Reactivity:

Human, Mouse

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, WB, IHC

Recommended dilutions:

ELISA:1:1000-1:2000, WB:1:200-1:1000,
IHC:1:25-1:100

Protein Background:

The protein encoded by this gene has both oxidoreductase and epimerase activities and is involved in androgen catabolism. The oxidoreductase activity can convert 3 alpha-adiol to dihydrotestosterone, while the epimerase activity can convert androsterone to epi-androsterone. Both reactions use NAD⁺ as the preferred cofactor. This gene is a member of the retinol dehydrogenase family.

Gene ID:

HSD17B6

Uniprot

O14756

Synonyms:

hydroxysteroid (17-beta) dehydrogenase 6


Immunogen:

Fusion protein of human HSD17B6.


Storage:

-20° C, pH7.4 PBS, 0.05% NaN₃, 40% Glycerol

Product Images



Gel: 10%SDS-PAGE, Lysate: 40 μ g, Lane: Mouse liver tissue, Primary antibody: PACO16495(HSD17B6 Antibody) at dilution 1/200, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 1 minute.



The image on the left is immunohistochemistry of paraffin-embedded Human colon cancer tissue using PACO16495(HSD17B6 Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification: x—200).