SSX2IP Antibody



PACO17154

Product Information

Size: Protein Background:

50ul This gene encodes a protein that binds the cancer-testis antigen Synovial Sarcoma X

Reactivity:

breakpoint 2 protein. The encoded protein may regulate the activity of Synovial

Reactivity:

Sarcoma V breakpoint 2 protein in malignant calls. Alternate collising results in malignant calls.

Sarcoma X breakpoint 2 protein in malignant cells. Alternate splicing results in multiple

Human transcript variants. A pseudogene of this gene is found on chromosome 3.

Source: Gene ID:

Rabbit SSX2IP

Isotype: Uniprot

IgG Q9Y2D8

Applications: Synonyms:

ELISA, WB, IHC synovial sarcoma, X breakpoint 2 interacting protein


Recommended dilutions: Immunogen:

ELISA:1:2000-1:5000, WB:1:500-1:2000, Fusion protein of human SSX2IP.

IHC:1:40-1:150 **Storage:**

-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product Images



The image on the left is immunohistochemistry of paraffin-embedded Human colon cancer tissue using PACO17154(SSX2IP Antibody) at dilution 1/45, on the right is treated with fusion protein. (Original magnification: x—200).

Gel: 8%SDS-PAGE, Lysate: 40 μ g, Lane: Hepg2 cells, Primary antibody: PACO17154(SSX2IP Antibody) at dilution 1/500, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 10 seconds.

The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using PACO17154(SSX2IP Antibody) at dilution 1/45, on the right is treated with fusion protein. (Original magnification: x—200).