
Product Information

Size:

50ul

Protein Background:

Vasohibin-2, also known as VASH2, is a 355 amino acid, protein belonging to the Vasohibin family. Expressed in various embryonic organs, Vasohibin-2 can be detected during embryonic weeks six through twelve and is found in vessels of 20-week embryonic organs as well as in endothelial cells of neonatal large vessels. Induced by VEGF, Vasohibin-2 is an angiogenesis inhibitor and interferes with the proliferation, migration and network formation by endothelial cells.

Reactivity:

Human, Mouse

Source:

Rabbit

Gene ID:**Isotype:**

IgG

VASH2

Uniprot**Applications:**

ELISA, WB

Q86V25

Recommended dilutions:

ELISA:1:2000-1:5000, WB:1:500-1:2000

vasohibin 2

Immunogen:


Full length fusion protein.

Synonyms:

-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Storage:

Product Images



Gel: 8%SDS-PAGE, Lysate: 40 μ g, Lane 1-2: HeLa and 293T cell, Primary antibody: PACO17415(VASH2 Antibody) at dilution 1/450 dilution, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 40 seconds.