AATK Antibody



PACO17473

Reactivity:

Human

Source:

Product Information

Size: Protein Background:

50ul The protein encoded by this gene contains a tyrosine kinase domain at the N-terminus

and a proline-rich domain at the C-terminus. This gene is induced during apoptosis, and expression of this gene may be a necessary pre-requisite for the induction of

growth arrest and/or apoptosis of myeloid precursor cells. This gene has been shown to produce neuronal differentiation in a neuroblastoma cell line. Two transcript variants

encoding different isoforms have been found for this gene.

Rabbit Gene ID:

Isotype: AATK

lgG Uniprot

Applications: Q6ZMQ8

ELISA, IHC Synonyms:

Recommended dilutions: Apoptosis-associated tyrosine kinase

ELISA:1:1000-1:2000, IHC:1:15-1:50 **Immunogen:**


Synthetic peptide of human AATK.

Storage:

-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product Images





The image on the left is immunohistochemistry of paraffin-embedded Human brain tissue using PACO17473(AATK Antibody) at dilution 1/10, on the right is treated with synthetic peptide. (Original magnification: x—200).

The image on the left is immunohistochemistry of paraffin-embedded Human breast cancer tissue using PACO17473(AATK Antibody) at dilution 1/10, on the right is treated with synthetic peptide. (Original magnification: x—200).