## **AGT Antibody**

PACO17530



| Product Information                  |   |
|--------------------------------------|---|
| Size:                                | Protein Background:   |
| 50ul                                 | The protein encoded by this gene, pre-angiotensinogen or angiotensinogen precursor,   |
| Reactivity:                          | is expressed in the liver and is cleaved by the enzyme renin in response to lowered blood pressure. The resulting product, angiotensin I, is then cleaved by angiotensin  |
| Human                                | converting enzyme (ACE) to generate the physiologically active enzyme angiotensin II.<br>The protein is involved in maintaining blood pressure and in the pathogenesis of |
| Source:                              | essential hypertension and preeclampsia. Mutations in this gene are associated with   |
| Rabbit                               | susceptibility to essential hypertension, and can cause renal tubular dysgenesis, a severe disorder of renal tubular development. Defects in this gene have also been     |
| lsotype:                             | associated with non-familial structural atrial fibrillation, and inflammatory bowel   |
| lgG                                  | disease.<br>Gene ID:  |
| Applications:                        | AGT   |
| ELISA, WB                            | Uniprot   |
| Recommended dilutions:               | P01019  |
| ELISA:1:1000-1:2000, WB:1:200-1:1000 | Synonyms:   |
|                                      | angiotensinogen (serpin peptidase inhibitor, clade A, member 8)   |
|                                      | Immunogen:  |
|                                      | Synthetic peptide of human AGT.   |
|                                      | Storage:  |
|                                      | -20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol   |

