
PRMT5 Antibody

PACO18364



Product Information	
Size:	Protein Background:
50ul	Protein arginine methyltransferase 5 has been shown to interact with WD repeat- containing protein 77, CLNS1A, Janus kinase 2, SNRPD3 and SUPT5H. Arginine methyltransferase that can both catalyze the formation of omega-N monomethylarginine (MMA) and symmetrical dimethylarginine (sDMA), with a preference for the formation of MMA. Specifically mediates the symmetrical dimethylation of arginine residues in the small nuclear ribonucleoproteins Sm D1 (SNRPD1) and Sm D3 (SNRPD3); such methylation being required for the assembly and biogenesis of snRNP core particles. Methylates SUPT5H. Mono- and dimethylates arginine residues of myelin basic protein (MBP) in vitro. Plays a role in the assembly of snRNP core particles. May play a role in cytokine-activated transduction pathways. Negatively regulates cyclin E1 promoter activity and cellular proliferation.
Reactivity:	
Human, Mouse, Rat	
Source:	
Rabbit	
lsotype:	
lgG	
Applications:	Gene ID:
ELISA, WB, IHC	PRMT5
Recommended dilutions:	Uniprot
ELISA:1:1000-1:2000, WB:1:200-1:1000, IHC:1:15-1:50	O14744
	Synonyms:
	protein arginine methyltransferase 5
	Immunogen:
	Synthetic peptide of human PRMT5.
	Storage:
	-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol



The image on the left is immunohistochemistry of paraffin-embedded Human tonsil tissue using PACO18364(PRMT5 Antibody) at dilution 1/20, on the right is treated with synthetic peptide. (Original magnification: x—200).

Gel: 10%SDS-PAGE, Lysate: 40 μ g, Lane 1-3: K562 cells, hela cells, 293T cells, Primary antibody: PACO18364(PRMT5 Antibody) at dilution 1/400, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 2 minutes.

The image on the left is immunohistochemistry of paraffin-embedded Human breast cancer tissue using PACO18364(PRMT5 Antibody) at dilution 1/20, on the right is treated with synthetic peptide. (Original magnification: x—200).