
PRKG2 Antibody

PACO18631



Product Information	
Size:	Protein Background:
50ul	Non-receptor protein-tyrosine kinase that plays an essential role in regulating cell migration, adhesion, spreading, reorganization of the actin cytoskeleton, formation and disassembly of focal adhesions and cell protrusions, cell cycle progression, cell proliferation and apoptosis. Required for early embryonic development and placenta development. Required for embryonic angiogenesis, normal cardiomyocyte migration and proliferation, and normal heart development. Regulates axon growth and neuronal cell migration, axon branching and synapse formation; required for normal development of the nervous system. Plays a role in osteogenesis and differentiation of osteoblasts. Functions in integrin signal transduction, but also in signaling downstream of numerous growth factor receptors, G-protein coupled receptors (GPCR), EPHA2, netrin receptors and LDL receptors.
Reactivity:	
Human, Mouse, Rat	
Source:	
Rabbit	
lsotype:	
lgG	
Applications:	Gene ID:
ELISA, WB, IHC	PRKG2
Recommended dilutions:	Uniprot
ELISA:1:2000-1:5000, WB:1:200-1:1000, IHC:1:25-1:100	Q13237
	Synonyms:
	protein kinase, cGMP-dependent, type II
	Immunogen:
	Synthetic peptide of human PRKG2.
	Storage:
	-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol



The image on the left is immunohistochemistry of paraffin-embedded Human cervical cancer tissue using PACO18631(PRKG2 Antibody) at dilution 1/20, on the right is treated with synthetic peptide. (Original magnification: x—200).

Gel: 8%SDS-PAGE, Lysate: 50 μ g, Lane 1-2: Mouse brain tissue, Mouse lung tissue, Primary antibody: PACO18631(PRKG2 Antibody) at dilution 1/160, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 30 seconds.

The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using PACO18631(PRKG2 Antibody) at dilution 1/20, on the right is treated with synthetic peptide. (Original magnification: x—200).