## **PAK3 Antibody**

## PACO18806





Size:	Protein Background:
50ul	G-protein coupled receptor for 5-hydroxytryptamine (serotonin). Also functions as a
Reactivity:	receptor for various drugs and psychoactive substances, including mescaline, psilocybin, 1-(2,5-dimethoxy-4-iodophenyl)-2-aminopropane (DOI) and lysergic acid,
Human, Mouse, Rat	diethylamide (LSD). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors. Beta-arrestin family members inhibit signaling via G
Source:	
Rabbit	proteins and mediate activation of alternative signaling pathways. Signaling activates phospholipase C and a phosphatidylinositol-calcium second messenger system that
lsotype:	modulates the activity of phosphatidylinositol 3-kinase and promotes the release of Ca2+ ions from intracellular stores. Affects neural activity, perception, cognition and
IgG	mood. Plays a role in the regulation of behavior, including responses to anxiogenic
Applications:	situations and psychoactive substances.
ELISA, WB, IHC	Gene ID:
Recommended dilutions:	PAK3
	Uniprot
ELISA:1:1000-1:2000, WB:1:200-1:1000, IHC:1:20-1:100	O75914
	Synonyms:
	p21 protein (Cdc42/Rac)-activated kinase 3
	Immunogen:
	Synthetic peptide of human PAK3.

## Storage:

-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol







The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using PACO18806(PAK3 Antibody) at dilution 1/25, on the right is treated with synthetic peptide. (Original magnification: x—200).

Gel: 6%SDS-PAGE, Lysate: 40 μ g, , Primary antibody: PACO18806(PAK3 Antibody) at dilution 1/200 dilution, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 1 minute.

The image on the left is immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using PACO18806(PAK3 Antibody) at dilution 1/25, on the right is treated with synthetic peptide. (Original magnification: x—200).