

### Product Information

**Size:**

50ul

**Reactivity:**

Human, Mouse, Rat

**Source:**

Rabbit

**Isotype:**

IgG

**Applications:**

ELISA, WB

**Recommended dilutions:**

ELISA:1:2000-1:5000, WB:1:500-1:2000

**Protein Background:**

A global transcription regulator. Complexes with cyclic AMP (cAMP) which allosterically activates DNA binding (to consensus sequence 5'-AAATGTGATCTAGATCACATTT-3') to directly regulate the transcription of about 300 genes in about 200 operons and indirectly regulate the expression of about half the genome. There are 3 classes of CRP promoters; class I promoters have a single CRP-binding site upstream of the RNA polymerase (RNAP)-binding site, whereas in class II promoters the single CRP- and RNAP-binding site overlap, CRP making multiple contacts with RNAP. Class III promoters require multiple activator molecules, including at least one CRP dimer. It can act as an activator, repressor, coactivator or corepressor. Induces a severe bend in DNA (about 87 degrees), bringing upstream promoter elements into contact with RNAP. Acts as a negative regulator of its own synthesis as well as for adenylate cyclase (*cyaA*), which generates cAMP. High levels of active CRP are detrimental to growth.

**Gene ID:**

CAV1

**Uniprot**

Q03135

**Synonyms:**

caveolin 1, caveolae protein, 22kDa

**Immunogen:**

Synthetic peptide of human CAV1.

**Storage:**

-20&deg; C, pH7.4 PBS, 0.05% NaN<sub>3</sub>, 40% Glycerol

## Product Images

---



Gel: 10%SDS-PAGE, Lysate: 40  $\mu$ g, Lane: Human leiomyosarcoma tissue, Primary antibody: PACO19257(CAV1 Antibody) at dilution 1/525, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 40 seconds.