## **PLIN2 Antibody**



## PACO19312

## **Product Information**

Size:

50ul

Reactivity:

Human, Mouse

Source:

Rabbit

Isotype:

lgG

**Applications:** 

ELISA, WB, IHC

Recommended dilutions:

ELISA:1:2000-1:5000, WB:1:500-1:2000, IHC:1:25-1:100

**Protein Background:** 

In muscle physiology, plays a central role in the balance between atrophy and hypertrophy. When recruited by MSTN, promotes atrophy response via phosphorylated SMAD2/4. MSTN decrease causes SMAD4 release and subsequent recruitment by the BMP pathway to promote hypertrophy via phosphorylated SMAD1/5/8. Acts synergistically with SMAD1 and YY1 in bone morphogenetic protein (BMP)-mediated cardiac-specific gene expression. Binds to SMAD binding elements (SBEs) (5'-GTCT/AGAC-3') within BMP response element (BMPRE) of cardiac activating regions. Common SMAD (co-SMAD) is the coactivator and mediator of signal transduction by TGF-beta (transforming growth factor). Component of the heterotrimeric SMAD2/SMAD3-SMAD4 complex that forms in the nucleus and is required for the TGF-mediated signaling. Promotes binding of the SMAD2/SMAD4/FAST-1 complex to DNA and provides an activation function required for SMAD1 or SMAD2 to stimulate transcription.

Gene ID:

PLIN2

Uniprot

Q99541

Synonyms:

perilipin 2


Immunogen:

Synthetic peptide of human PLIN2.


Storage:

-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol


## **Product Images**



The image on the left is immunohistochemistry of paraffin-embedded Human gastric cancer tissue using PACO19312(PLIN2 Antibody) at dilution 1/30, on the right is treated with synthetic peptide. (Original magnification: x—200).



Gel: 8%SDS-PAGE, Lysate: 40 μ g, Lane: Mouse brain tissue, Primary antibody: PACO19312(PLIN2 Antibody) at dilution 1/680, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 1 minute.



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using PACO19312(PLIN2 Antibody) at dilution 1/30, on the right is treated with synthetic peptide. (Original magnification: x—200).