GCGR Antibody

PACO19714




Size:	Protein Background:
50ul	Histone demethylase that demethylates both 'Lys-4' (H3K4me) and 'Lys-9' (H3K9me) of
Reactivity:	histone H3, thereby acting as a coactivator or a corepressor, depending on the context. Acts by oxidizing the substrate by FAD to generate the corresponding imine that is
Human, Mouse, Rat	subsequently hydrolyzed. Acts as a corepressor by mediating demethylation of H3K4me, a specific tag for epigenetic transcriptional activation. Demethylates both
Source:	mono- (H3K4me1) and di-methylated (H3K4me2) H3K4me. May play a role in the
Rabbit	repression of neuronal genes. Alone, it is unable to demethylate H3K4me on nucleosomes and requires the presence of RCOR1/CoREST to achieve such activity. Also
lsotype:	acts as a coactivator of androgen receptor (ANDR)-dependent transcription, by being recruited to ANDR target genes and mediating demethylation of H3K9me, a specific
lgG	tag for epigenetic transcriptional repression.
Applications:	Gene ID:
ELISA, IHC	GCGR
Recommended dilutions:	Uniprot
ELISA:1:1000-1:2000, IHC:1:25-1:100	P47871
	Synonyms:
	glucagon receptor
	glucagon receptor Immunogen:

Storage:

-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol





The image on the left is immunohistochemistry of paraffin-embedded Human brain tissue using PACO19714(GCGR Antibody) at dilution 1/25, on the right is treated with synthetic peptide. (Original magnification: x—200).