# **HRG Antibody**

**Product Information** 



#### PACO19777

## Size: **Protein Background:** 50ul Functions in cell-cell adhesion, cell migration and axon guidance, exerting an attractive or repulsive role depending on its interaction partners. Plays a role in the spatial Reactivity: organization of brain neurons. Plays a role in vascular development in the retina. Plays a role in cell-cell adhesion via its interaction with ADGRL3 and probably also other Human latrophilins that are expressed at the surface of adjacent cells. Interaction with the Source: intracellular domain of ROBO1 mediates axon attraction towards cells expressing NTN1. Mediates axon growth cone collapse and plays a repulsive role in neuron guidance via Rabbit its interaction with UNC5B, and possibly also other UNC-5 family members. Promotes neurite outgrowth (in vitro). Mediates cell-cell contacts that promote an increase both Isotype: in neurite number and in neurite length. Plays a role in the regulation of the density of lgG glutamaergic synapses. Plays a role in fibroblast growth factor-mediated signaling cascades. **Applications:** Gene ID: ELISA, IHC HRG **Recommended dilutions:** Uniprot ELISA:1:2000-1:5000, IHC:1:50-1:200 P04196 Synonyms: histidine-rich glycoprotein


### Storage:


Immunogen:

-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Synthetic peptide of human HRG.

# **Product Images**





The image on the left is immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using PACO19777(HRG Antibody) at dilution 1/50, on the right is treated with synthetic peptide. (Original magnification: x—200).

The image on the left is immunohistochemistry of paraffin-embedded Human breast cancer tissue using PACO19777(HRG Antibody) at dilution 1/50, on the right is treated with synthetic peptide. (Original magnification: x—200).