SCD Antibody



PACO20400

Product Information

Size:

50ul

Reactivity:

Human

Source:

Rabbit

Isotype:

lgG

Applications:

ELISA, WB, IHC

Recommended dilutions:

ELISA:1:1000-1:2000, WB:1:200-1:1000, IHC:1:10-1:50

Protein Background:

Plays a role in vesicle-mediated protein trafficking to lysosomal compartments including the endocytic membrane transport and autophagic pathways. Believed to act as a core component of the putative HOPS and CORVET endosomal tethering complexes which are proposed to be involved in the Rab5-to-Rab7 endosome conversion probably implicating MON1A/B, and via binding SNAREs and SNARE complexes to mediate tethering and docking events during SNARE-mediated membrane fusion. The HOPS complex is proposed to be recruited to Rab7 on the late endosomal membrane and to regulate late endocytic, phagocytic and autophagic traffic towards lysosomes. The CORVET complex is proposed to function as a Rab5 effector to mediate early endosome fusion probably in specific endosome subpopulations. Required for recruitment of VPS33A to the HOPS complex. Required for fusion of endosomes and autophagosomes with lysosomes; the function is dependent on its association with VPS33A but not VPS33B.

Gene ID:

SCD

Uniprot

O00767

Synonyms:

stearoyl-CoA desaturase (delta-9-desaturase)


Immunogen:


Synthetic peptide of human SCD.


Storage:


-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol


Product Images











The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using PACO20400(SCD Antibody) at dilution 1/20, on the right is treated with synthetic peptide. (Original magnification: x—200).