UCP3 Antibody



PACO20819

Product Information

Size:

50ul

Reactivity:

Human, Mouse, Rat

Source:

Rabbit

Isotype:

lgG

Applications:

ELISA, WB, IHC

Recommended dilutions:

ELISA:1:1000-1:2000, WB:1:200-1:1000, IHC:1:10-1:50

Protein Background:

Transcriptional repressor. Recognizes and binds to the consensus sequence '5-[CG]NG[CG]GGCA[CA]CC-3'. May act as a tumor suppressor. May be involved in development of head, face, limbs and ventral body wall. Involved in down-regulation of SIRT1 and thereby is involved in regulation of p53/TP53-dependent apoptotic DNA-damage responses. The specific target gene promoter association seems to be depend on corepressors, such as CTBP1 or CTBP2 and MTA1. The regulation of SIRT1 transcription in response to nutrient deprivation seems to involve CTBP1. In cooperation with MTA1 (indicative for an association with the NuRD complex) represses transcription from CCND1/cyclin-D1 and CDKN1C/p57Kip2 specifically in quiescent cells. Involved in regulation of the Wnt signaling pathway probably by association with TCF7L2 and preventing TCF7L2 and CTNNB1 association with promoters of TCF-responsive genes.

Gene ID:

UCP3

Uniprot

P55916

Synonyms:

uncoupling protein 3 (mitochondrial, proton carrier)


Immunogen:


Synthetic peptide of human UCP3.


Storage:

-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol


Product Images







Gel: 8%SDS-PAGE, Lysate: 40 μ gPrimary antibody: PACO20819(UCP3 Antibody) at dilution 1/200 dilution, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 1 minute.



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using PACO20819(UCP3 Antibody) at dilution 1/20, on the right is treated with synthetic peptide. (Original magnification: x—200).