## **WNT8A Antibody**



## PACO20919


## **Product Information** Size: **Protein Background:** 50ul Originally identified as neuronal protein that specifically associates with HTT/huntingtin and the binding is enhanced by an expanded polyglutamine repeat within HTT possibly Reactivity: affecting HAP1 interaction properties. Both HTT and HAP1 are involved in intracellular trafficking and HAP1 is proposed to link HTT to motor proteins and/or transport cargos. Human Seems to play a role in vesicular transport within neurons and axons such as from early Source: endosomes to late endocytic compartments and to promote neurite outgrowth. The vesicular transport function via association with microtubule-dependent transporters Rabbit can be attenuated by association with mutant HTT. Involved in the axonal transport of BDNF and its activity-dependent secretion; the function seems to involve HTT, DCTN1 Isotype: and a complex with SORT1. Involved in APP trafficking and seems to faciltate APP lgG anterograde transport and membrane insertion thereby possibly reducing processing into amyloid beta. **Applications:** Gene ID: ELISA, IHC WNT8A **Recommended dilutions:** Uniprot ELISA:1:2000-1:5000, IHC:1:25-1:100 Q9H1J5 Synonyms: wingless-type MMTV integration site family, member 8A Immunogen:

-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Synthetic peptide of human WNT8A.

Storage:

## **Product Images**



The image on the left is immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using PACO20919(WNT8A Antibody) at dilution 1/20, on the right is treated with synthetic peptide. (Original magnification: x—200).