BAZ1B Antibody

PACO20927



Product Information	
Size:	Protein Background:
50ul	Functions in nuclear protein import, either in association with an adapter protein, like an importin-alpha subunit, which binds to nuclear localization signals (NLS) in cargo substrates, or by acting as autonomous nuclear transport receptor. Acting autonomously, serves itself as NLS receptor. Docking of the importin/substrate complex to the nuclear pore complex (NPC) is mediated by KPNB1 through binding to nucleoporin FxFG repeats and the complex is subsequently translocated through the pore by an energy requiring, Ran-dependent mechanism. At the nucleoplasmic side of the NPC, Ran binds to importin-beta and the three components separate and importinalpha and -beta are re-exported from the nucleus to the cytoplasm where GTP hydrolysis releases Ran from importin. The directionality of nuclear import is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus. Mediates autonomously the nuclear import of ribosomal proteins RPL23A, RPS7 and RPL5.
Reactivity:	
Human, Mouse	
Source:	
Rabbit	
lsotype:	
lgG	
Applications:	
ELISA, IHC	Gene ID:
Recommended dilutions:	BAZ1B
ELISA:1:2000-1:5000, IHC:1:25-1:100	Uniprot
	Q9UIG0
	Synonyms:
	bromodomain adjacent to zinc finger domain, 1B
	Immunogen:
	Synthetic peptide of human BAZ1B.
	Storage:

-20° C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol



The image on the left is immunohistochemistry of paraffin-embedded Human breast cancer tissue using PACO20927(BAZ1B Antibody) at dilution 1/35, on the right is treated with synthetic peptide. (Original magnification: x—200).