FAS Antibody



PACO23142

Reactivity:

IHC:1:50-1:100

Human

Product Information

Size: **Protein Background:**

100ul Receptor for TNFSF6/FASLG. The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs

> caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. FAS-mediated apoptosis

may have a role in the induction of peripheral tolerance, in the antigen-stimulated Source:

suicide of mature T-cells, or both. The secreted isoforms 2 to 6 block apoptosis (in vitro).

Rabbit

Gene ID: Isotype:

FAS lgG

Uniprot **Applications:**

P25445 ELISA, WB, IHC

Synonyms: **Recommended dilutions:**


Apo-1; CD95; FasL receptor; TNFRSF6; TNR6 ELISA:1:2000-1:10000, WB:1:500-1:3000,

Immunogen: Synthesized peptide derived from C-terminal of human Fas.


Storage:

Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Product Images



Western blot analysis of extracts from LOVO cells, using Fas antibody.



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue using Fas antibody.