

Product Information

Size:

100ul

Reactivity:

Human, Mouse, Rat

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, WB, IHC

Recommended dilutions:

ELISA:1:2000-1:10000, WB:1:500-1:3000,
IHC:1:50-1:100

Protein Background:

Functions in mitochondrial tRNA maturation. Part of mitochondrial ribonuclease P, an enzyme composed of MRPP1/TRMT10C, MRPP2/HSD17B10 and MRPP3/KIAA0391, which cleaves tRNA molecules in their 5'-ends. Catalyzes the beta-oxidation at position 17 of androgens and estrogens and has 3-alpha-hydroxysteroid dehydrogenase activity with androsterone. Catalyzes the third step in the beta-oxidation of fatty acids. Carries out oxidative conversions of 7-alpha-OH and 7-beta-OH bile acids. Also exhibits 20-beta-OH and 21-OH dehydrogenase activities with C21 steroids. By interacting with intracellular amyloid-beta, it may contribute to the neuronal dysfunction associated with Alzheimer disease (AD).

Gene ID:

HSD17B10

Uniprot

Q99714

Synonyms:

EC 1.1.1.35; 3-hydroxyacyl-CoA dehydrogenase type II; Type II HADH; 3-hydroxy-2-methylbutyryl-CoA dehydrogenase; EC 1.1.1.178

Immunogen:


Synthesized peptide derived from human ERAB.

Storage:


Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Product Images

Immunohistochemical analysis of paraffin-embedded human brain tissue using ERAB antibody.



Western blot analysis of extracts from A549 cells (Lane 2) and HeLa cells (Lane 3), using ERAB antibody. The lane on the left is treated with synthesized peptide.



Western blot analysis of extracts from LOVO cells, using ERAB antibody.

