ASAH2 Antibody, Biotin conjugated

PACO25079



Product Information	
Size:	Protein Background:
50ug	Hydrolyzes the sphingolipid ceramide into sphingosine and free fatty acid, at an
Reactivity:	optimal pH of 6.5-8.5. Acts as a key regulator of sphingolipid signaling metabolites by generating sphingosine at the cell surface. Acts as a repressor of apoptosis both by
Human	reducing C16-ceramide, thereby preventing ceramide-induced apoptosis, and generating sphingosine, a precursor of the antiapoptotic factor sphingosine 1-
Source:	phosphate. Probably involved in the digestion of dietary sphingolipids in intestine by
Rabbit	acting as a key enzyme for the catabolism of dietary sphingolipids and regulating the levels of bioactive sphingolipid metabolites in the intestinal tract.
lsotype:	Gene ID:
lgG	ASAH2
Applications:	Uniprot
ELISA	Q9NR71
Recommended dilutions:	Synonyms:
	Neutral ceramidase (N-CDase) (NCDase) (EC 3.5.1.23) (Acylsphingosine deacylase 2) (BCDase) (LCDase) (hCD) (N-acylsphingosine amidohydrolase 2) (Non-lysosomal ceramidase) [Cleaved into: Neutral ceramidase soluble form], ASAH2, HNAC1
	Immunogen:
	Recombinant Human Neutral ceramidase protein (255-506AA).
	Storage:
	Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

N/A N/A