

PACO27341

Product Information

Size:

50ug

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, IHC, IF

Recommended dilutions:

ELISA:1:2000-1:10000, IHC:1:20-1:200,
IF:1:50-1:200

Protein Background:

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Also involved in the deacetylation of cohesin complex protein SMC3 regulating release of cohesin complexes from chromatin. May play a role in smooth muscle cell contractility.

Gene ID:

HDAC8

Uniprot

Q9BY41

Synonyms:

Histone deacetylase 8 (HD8) (EC 3.5.1.98), HDAC8, HDACL1


Immunogen:

Recombinant Human Histone deacetylase 8 protein (1-377AA).


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4


Product Images



Immunohistochemistry of paraffin-embedded human placenta tissue using PACO27341 at dilution of 1:100.



Immunofluorescent analysis of HeLa cells using PACO27341 at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemistry of paraffin-embedded human breast cancer using PACO27341 at dilution of 1:100.