CDK4 Antibody, HRP conjugated

PACO33093



Product Information	
Size:	Protein Background:
50ug	Ser/Thr-kinase component of cyclin D-CDK4 (DC) complexes that phosphorylate and
Reactivity:	inhibit members of the retinoblastoma (RB) protein family including RB1 and regulate the cell-cycle during G1/S transition. Phosphorylation of RB1 allows dissociation of the
Human	transcription factor E2F from the RB/E2F complexes and the subsequent transcription of E2F target genes which are responsible for the progression through the G1 phase.
Source:	Hypophosphorylates RB1 in early G1 phase. Cyclin D-CDK4 complexes are major
Rabbit	integrators of various mitogenenic and antimitogenic signals. Also phosphorylates SMAD3 in a cell-cycle-dependent manner and represses its transcriptional activity.
lsotype:	Component of the ternary complex, cyclin D/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex.
lgG	Gene ID:
Applications:	CDK4
ELISA	Uniprot
Recommended dilutions:	P11802
	Synonyms:
	Cyclin-dependent kinase 4 (EC 2.7.11.22) (Cell division protein kinase 4) (PSK-J3), CDK4
	Immunogen:
	Recombinant Human Cyclin-dependent kinase 4 protein (2-303AA).
	Storage:
	Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

N/A N/A