## HA Antibody

PACO33964



Product Information	
Size:	Protein Background:
50ug	Binds to sialic acid, containing receptors on the cell surface, bringing about the
Reactivity:	attachment of the virus particle to the cell. This attachment induces virion internalization of about two third of the virus particles through clathrin-dependent
Influenza A virus(H7N9)	endocytosis and about one third through a clathrin- and caveolin-independent pathway. Plays a major role in the determination of host range restriction and virulence.
Source:	Class I viral fusion protein. Responsible for penetration of the virus into the cell
Rabbit	cytoplasm by mediating the fusion of the membrane of the endocytosed virus particle with the endosomal membrane. Low pH in endosomes induces an irreversible
lsotype:	conformational change in HA2, releasing the fusion hydrophobic peptide. Several
lgG	trimers are required to form a competent fusion pore.
Applications:	HA
ELISA	Uniprot
Recommended dilutions:	R4NN21
	Synonyms:
	Hemagglutinin [Cleaved into: Hemagglutinin HA1 chain; Hemagglutinin HA2 chain], HA
	Immunogen:
	Recombinant Influenza A virus Hemagglutinin protein (19-524AA).
	Storage:
	Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

N/A N/A